



WITH

*nutricote* GAL-Xe<sup>ONE</sup>

# 17-4-8 NPK PLUS

**GUARANTEED ANALYSIS:**

|  |            |
|--|------------|
| <b>Total Nitrogen (N)*</b> .....                               | <b>17%</b> |
| 8.40% Nitrate Nitrogen   |            |
| 8.60% Ammoniacal Nitrogen                                      |            |
| <b>Available Phosphate (P<sub>2</sub>O<sub>5</sub>)*</b> ..... | <b>4%</b>  |
| <b>Soluble Potash (K<sub>2</sub>O)*</b> .....                  | <b>8%</b>  |
| Magnesium (Mg)*.....   | 1.30%      |
| 0.80% Water Soluble Magnesium (Mg)                             |            |
| Sulfur (S).....  | 2.40%      |
| 2.40% Combined Sulfur (S)                                      |            |
| Boron (B)*.....  | 0.02%      |
| Copper (Cu)*.....  | 0.05%      |
| 0.02% Water Soluble Copper (Cu)                                |            |
| Iron (Fe).....   | 0.99%      |
| 0.17% Chelated Iron (Fe)                                       |            |
| 0.27% Water Soluble Iron (Fe)                                  |            |
| Manganese (Mn)*.....   | 0.09%      |
| 0.03% Water Soluble Manganese (Mn)                             |            |
| Molybdenum (Mo)*.....  | 0.005%     |

**DERIVED FROM:** Polymer Coated: Ammonium Nitrate, Ammonium Phosphate, Calcium Phosphate, Ammonium Polyphosphate, Potassium Sulfate, Magnesium Sulfate, Iron EDTA, Sodium Borate, Copper Sulfate, Manganese Sulfate, Sodium Molybdate; Ferrous Sulfate, Iron Sucrate, Iron Citrate, Magnesium Sulfate, Magnesium Sucrate, Manganese Sucrate, Copper Sucrate, Sodium Borate, Sodium Molybdate | \*The Ammonium Nitrate, Ammonium Phosphate, Calcium Phosphate, Ammonium Polyphosphate, Potassium Sulfate, Magnesium Sulfate, Sodium Borate, Copper Sulfate, Manganese Sulfate, and Sodium Molybdate have been polymer coated to provide 17% Slow Release Nitrogen (N), 4% Slow Release Available Phosphate (P<sub>2</sub>O<sub>5</sub>), 8% Slow Release Soluble Potash (K<sub>2</sub>O), 0.10% Slow Release Iron (Fe), 0.6% Slow Release Magnesium (Mg), 0.01% Slow Release Boron (B), 0.02% Slow Release Copper (Cu), 0.03% Slow Release Manganese (Mn), and 0.005% Slow Release Molybdenum (Mo).

The below rates are provided as a guide only. Specific rates will vary by region, soil type, and performance requirements. Before deciding which formulation, release type, and rate to use, contact your Florikan Technical Representative. They will advise you in choosing the right product for your crop and conditions.

| INCORPORATION RATES - LBS PER CUBIC YARD |     |     |      |       |
|--|-----|-----|------|-------|
|  | LOW | MED | HIGH | HEAVY |
| LBS                                      | 6   | 10  | 14   | 17    |

  

| BROADCAST RATES - LBS PER 1000SQFT |     |     |      |       |
|------------------------------------|-----|-----|------|-------|
|                                    | LOW | MED | HIGH | HEAVY |
| LBS                                | 8   | 10  | 11   | n/a   |

**NOTE:** low/med/high is crop specific information. HIGH refers to heavy feeding crop, LOW is for sensitive feeding crop.

**Conversion**

5 Grams = 1 Teaspoon | 15 Grams = 1 Tablespoon

| TOP DRESS RATES - GRAMS PER CONTAINER |     |     |      |       |
|---------------------------------------|-----|-----|------|-------|
| VOL                                   | LOW | MED | HIGH | HEAVY |
| 1 GAL                                 | 10  | 15  | 20   | 30    |
| 3 GAL                                 | 30  | 45  | 60   | 75    |
| 7 GAL                                 | 50  | 60  | 90   | 110   |
| 10 GAL                                | 60  | 90  | 135  | 160   |
| 15 GAL                                | 75  | 125 | 150  | 190   |
| 25 GAL                                | 125 | 175 | 250  | 300   |
| 45 GAL                                | 200 | 275 | 450  | 540   |
| 65 GAL                                | 300 | 425 | 650  | 685   |

**SKU: 17-4-8-180**

**U.S. STANDARD - NET WEIGHT 50 LB. (22.68 KG.)**



This product was developed in conjunction; and with assistance from NASA using cutting edge research, quality control and scientific methodology.



**Signal Word: WARNING**

**Hazard Statement:** Causes skin and eye irritation - may cause respiratory irritation

**Precautionary Statement:** Do not inhale - handle in well-ventilated areas - wear protective gloves - wash thoroughly after handling - if inhaled, move victim to fresh air and in comfortable breathing position - call doctor/poison control if swallowed/inhaled, skin/eyes become irritated, or feel unwell - wash contaminated clothes before reuse - store locked up & in well ventilated area - keep container tightly closed - dispose per US laws



813430010637

**UP TO:  
180 DAYS**

**AVG MEDIA TEMPS**

| 60°        |   | 70°        |   | 80°        |   |
|------------|---|------------|---|------------|---|
| 240        | 8 | 210        | 7 | 180        | 6 |
| Days/Mnths |   | Days/Mnths |   | Days/Mnths |   |

**GUARANTEED BY: F1062**

**QUALITY CONTROL BATCH NUMBER:**



**WARNING TO PURCHASER, THIS IS A DISCLAIMER.** The performance warranty and/or release of this product, whether stated or implied, is null and void if improperly stored.

**Heavy Metal Statement:** Information regarding the contents and levels of metals in this product is available on the internet at <http://www.aapfco.org/metals.html>

**CAUTION:** KEEP OUT OF THE REACH OF CHILDREN HARMFUL IF SWALLOWED. This product must be stored under cool and dry conditions.

**Environmental Caution:** As a precaution, fertilizer products should not be applied within 10 feet of any water body.

**Caution:** The application of fertilizer containing Molybdenum (Mo) may result in forage crops containing levels of Molybdenum which may be toxic to ruminant animals.

**WARNING:** This product contains Boron (B) and should be used according to directions only, especially on crops sensitive to Boron (B) uptake.

**NOTE:** Fertilizer products can be hygroscopic and may be affected by humidity and/or heat during improper storage. Please keep this fertilizer product in a dry condition and away from heat and moisture which may start release in the bag.